

Fixed-point elimination in Heyting algebras¹

Luigi Santocanale
LIF, Aix-Marseille Université

SSAOS, Trojanovice, September 2016

Work in collaboration with
Silvio Ghilardi (U. of Milan) and Maria João Gouveia (U. of Lisbon)

¹See [Ghilardi et al., 2016]

Plan

A primer on mu-calculi

The intuitionistic μ -calculus

The elimination procedure

Bounding closure ordinals

Add to a given algebraic framework
syntactic least and greatest fixed-point constructors.

μ -calculi

Add to a given algebraic framework
syntactic least and greatest fixed-point constructors.

E.g., the propositional modal μ -calculus:

$$\begin{aligned} \phi := & x \mid \neg x \mid \top \mid \phi \wedge \phi \mid \perp \mid \phi \vee \phi \mid \Box \phi \mid \Diamond \phi \\ & \mid \mu_x.\phi \mid \nu_x.\phi, \quad \text{when } x \text{ is positive in } \phi. \end{aligned}$$

μ -calculi

Add to a given algebraic framework
syntactic least and greatest fixed-point constructors.

E.g., the propositional modal μ -calculus:

$$\begin{aligned} \phi := & x \mid \neg x \mid \top \mid \phi \wedge \phi \mid \perp \mid \phi \vee \phi \mid \Box \phi \mid \Diamond \phi \\ & \mid \mu_x.\phi \mid \nu_x.\phi, \quad \text{when } x \text{ is positive in } \phi. \end{aligned}$$

Interpret the syntactic least (resp. greatest) fixed-point as expected.

$$\begin{aligned} \llbracket \mu_x.\phi \rrbracket_v := \\ \text{least fixed-point of the monotone mapping } X \mapsto \llbracket \phi \rrbracket_{v, X/x} \end{aligned}$$

Alternation hierarchies in μ -calculi

Let $\#$ count the number of alternating blocks of fixed-points.

Alternation hierarchies in μ -calculi

Let $\#$ count the number of alternating blocks of fixed-points.

Problem. For a given μ -calculus, does there exist n such that, for each ϕ with $\#\phi > n$, there exists ψ with $\gamma \equiv \psi$ and $\#\psi \leq n$?

Alternation hierarchies: fatcs

- ▶ The alternation hierarchy for the modal μ -calculus is infinite (there exists no such n) [Lenzi, 1996, Bradfield, 1998].
- ▶ Idem for the lattice μ -calculus [Santocanale, 2002].
- ▶ The alternation hierarchy for the linear μ -calculus ($\diamond x = \square x$) is reduced to the Büchi fragment (here $n = 2$).
- ▶ The alternation hierarchy for the modal μ -calculus on transitive frames collapses to the alternation free fragment (here $n = 1.5$) [Alberucci and Facchini, 2009].
- ▶ The alternation hierarchy for the distributive μ -calculus is trivial (here $n = 0$) [Kozen, 1983].

Alternation hierarchies: fatcs

- ▶ The alternation hierarchy for the modal μ -calculus is infinite (there exists no such n) [Lenzi, 1996, Bradfield, 1998].
- ▶ Idem for the lattice μ -calculus [Santocanale, 2002].
- ▶ The alternation hierarchy for the linear μ -calculus ($\diamond x = \square x$) is reduced to the Büchi fragment (here $n = 2$).
- ▶ The alternation hierarchy for the modal μ -calculus on transitive frames collapses to the alternation free fragment (here $n = 1.5$) [Alberucci and Facchini, 2009].
- ▶ The alternation hierarchy for the distributive μ -calculus is trivial (here $n = 0$) [Kozen, 1983].

A research plan:

Develop a theory explaining why alternation hierarchies collapses.

μ -calculi on generalized distributive lattices

Theorem. [Frittella and Santocanale, 2014] There are lattice varieties (Nation's varieties)

$$\mathcal{D}_0 \subseteq \mathcal{D}_1 \subseteq \dots \subseteq \mathcal{D}_n \subseteq \dots$$

with \mathcal{D}_0 the variety of distributive lattices, such that, on \mathcal{D}_n and for any lattice term ϕ ,

$$\phi^{n+2}(\perp) = \phi^{n+1}(\perp) \quad (= \mu_x.\phi), \quad \phi^n(\perp) \neq \phi^{n+1}(\perp).$$

μ -calculi on generalized distributive lattices

Theorem. [Frittella and Santocanale, 2014] There are lattice varieties (Nation's varieties)

$$\mathcal{D}_0 \subseteq \mathcal{D}_1 \subseteq \dots \subseteq \mathcal{D}_n \subseteq \dots$$

with \mathcal{D}_0 the variety of distributive lattices, such that, on \mathcal{D}_n and for any lattice term ϕ ,

$$\phi^{n+2}(\perp) = \phi^{n+1}(\perp) \quad (= \mu_x.\phi), \quad \phi^n(\perp) \neq \phi^{n+1}(\perp).$$

Corollary. The alternation hierarchy of the lattice μ -calculus is trivial on \mathcal{D}_n , for each $n \geq 0$.

Plan

A primer on mu-calculi

The intuitionistic μ -calculus

The elimination procedure

Bounding closure ordinals

The intuitionistic μ -calculus

After the distributive μ -calculus, the next on the list.

We extend the signature of Heyting algebras (i.e. Intuitionistic Logic) with least and greatest fixed-point constructors.

Intuitionistic μ -terms are generated by the grammar:

$$\begin{aligned} \phi := x \mid \top \mid \phi \wedge \phi \mid \perp \mid \phi \vee \phi \mid \phi \rightarrow \phi \\ \mid \mu_x.\phi \mid \nu_x.\phi, \quad \text{when } x \text{ is positive in } \phi. \end{aligned}$$

Semantics

We take any *provability* semantics of IL with fixed points:

- ▶ (Complete) Heyting algebras.
- ▶ Kripke frames.
- ▶ Any sequent calculus for Intuitionistic Logic (e.g. LJ) plus Park/Kozen's rules for least and greatest fixed-points:

$$\frac{\phi[\psi/x] \vdash \psi}{\mu_x.\phi \dashv \psi} \qquad \frac{\Gamma \vdash \phi(\mu_x.\phi)}{\Gamma \vdash \mu_x.\phi}$$

$$\frac{\phi(\nu_x.\phi) \vdash \delta}{\nu_x.\phi \vdash \delta} \qquad \frac{\psi \vdash \phi[\psi/x]}{\psi \vdash \nu_x.\phi}$$

Heyting algebras

Definition. A *Heyting algebra* is a bounded lattice $H = \langle H, \top, \wedge, \perp, \vee \rangle$ with an additional binary operation \rightarrow satisfying

$$x \wedge y \leq z \quad \text{iff} \quad x \leq y \rightarrow z.$$

Pitt's (bisimulation) quantifiers

Proposition. [Pitts, 1992, Ghilardi and Zawadowski, 1997] Given ϕ with $x \in \text{Var}(\phi)$,

we can construct a term $\exists_x.\phi$

such that

1. $\text{Var}(\exists_x.\phi) \subseteq \text{Var}(\phi) \setminus \{x\}$, and
2. whenever $\text{Var}(\psi) \subseteq \text{Var}(\phi) \setminus \{x\}$, we have

$$\phi \leq \psi \quad \text{iff} \quad \exists_x.\phi \leq \psi.$$

Pitt's (bisimulation) quantifiers

Proposition. [Pitts, 1992, Ghilardi and Zawadowski, 1997] Given ϕ with $x \in \text{Var}(\phi)$,

we can construct a term $\exists_x.\phi$

such that

1. $\text{Var}(\exists_x.\phi) \subseteq \text{Var}(\phi) \setminus \{x\}$, and
2. whenever $\text{Var}(\psi) \subseteq \text{Var}(\phi) \setminus \{x\}$, we have

$$\phi \leq \psi \quad \text{iff} \quad \exists_x.\phi \leq \psi.$$

NB: we can also construct $\forall_x.\phi$.

Pitt's (bisimulation) quantifiers

Proposition. [Pitts, 1992, Ghilardi and Zawadowski, 1997] Given ϕ with $x \in \text{Var}(\phi)$,

we can construct a term $\exists_x.\phi$

such that

1. $\text{Var}(\exists_x.\phi) \subseteq \text{Var}(\phi) \setminus \{x\}$, and
2. whenever $\text{Var}(\psi) \subseteq \text{Var}(\phi) \setminus \{x\}$, we have

$$\phi \leq \psi \quad \text{iff} \quad \exists_x.\phi \leq \psi.$$

NB: we can also construct $\forall_x.\phi$.

Define least and greatest fixed points from the quantifiers.

Ruitenburg's theorem [Ruitenburg, 1984]

Theorem. For each intuitionistic formula ϕ , there exists $n \geq 0$ such that $\phi^n(x) \equiv \phi^{n+2}(x)$.

Ruitenburg's theorem [Ruitenburg, 1984]

Theorem. For each intuitionistic formula ϕ , there exists $n \geq 0$ such that $\phi^n(x) \equiv \phi^{n+2}(x)$.

Then

$$\phi^n(\perp) \leq \phi^{n+1}(\perp) \leq \phi^{n+2}(\perp) = \phi^n(\perp),$$

so $\phi^n(\perp)$ is the least fixed-point of ϕ .

Ruitenburg's theorem [Ruitenburg, 1984]

Theorem. For each intuitionistic formula ϕ , there exists $n \geq 0$ such that $\phi^n(x) \equiv \phi^{n+2}(x)$.

Then

$$\phi^n(\perp) \leq \phi^{n+1}(\perp) \leq \phi^{n+2}(\perp) = \phi^n(\perp),$$

so $\phi^n(\perp)$ is the least fixed-point of ϕ .

Corollary. The alternation hierarchy for the intuitionistic μ -calculus is trivial.

Ruitenburg's theorem [Ruitenburg, 1984]

Theorem. For each intuitionistic formula ϕ , there exists $n \geq 0$ such that $\phi^n(x) \equiv \phi^{n+2}(x)$.

Then

$$\phi^n(\perp) \leq \phi^{n+1}(\perp) \leq \phi^{n+2}(\perp) = \phi^n(\perp),$$

so $\phi^n(\perp)$ is the least fixed-point of ϕ .

Corollary. The alternation hierarchy for the intuitionistic μ -calculus is trivial.

NB : Ruitenburg's n might not be the *closure ordinal* of $\mu_x.\phi$.

Peirce, compatibility, strengths and strongness

Proposition. Peirce's theorem for Heyting algebras.

Every term ϕ is compatible. In particular, for ψ, χ arbitrary terms, the equation

$$\phi[\psi/x] \wedge \chi = \phi[\psi \wedge \chi/x] \wedge \chi.$$

holds on Heyting algebras.

Corollary. Every term ϕ monotone in x is *strong* in x . That is, any the following equivalent conditions

$$\phi[\psi/x] \wedge \chi \leq \phi[\psi \wedge \chi/x], \quad \psi \rightarrow \chi \leq \phi[\psi/x] \rightarrow \phi[\chi/x],$$

hold, for any terms ψ and χ .

Plan

A primer on mu-calculi

The intuitionistic μ -calculus

The elimination procedure

Bounding closure ordinals

Greatest fixed-points

Proposition. On Heyting algebras, we have

$$\nu_x.\phi = \phi(\top).$$

Greatest fixed-points

Proposition. On Heyting algebras, we have

$$\nu_x.\phi = \phi(\top).$$

The deduction theorem and Pitts quantifiers imply that

$$\nu_x.\phi(x) = \exists_x.(x \wedge x \rightarrow \phi(x)) = \phi(\top).$$

Greatest fixed-points

Proposition. On Heyting algebras, we have

$$\nu_x.\phi = \phi(\top).$$

The deduction theorem and Pitts quantifiers imply that

$$\nu_x.\phi(x) = \exists_x.(x \wedge x \rightarrow \phi(x)) = \phi(\top).$$

Using strongness:

$$\phi(\top) = \phi(\top) \wedge \phi(\top) \leq \phi(\top \wedge \phi(\top)) = \phi^2(\top).$$

Greatest solutions of systems of equations

Proposition. On Heyting algebras, a system of equations

$$\left\{ \begin{array}{l} x_1 = \phi_1(x_1, \dots, x_n) \\ \vdots \\ x_n = \phi_n(x_1, \dots, x_n) \end{array} \right\}$$

has a greatest solution obtained by iterating

$$\phi := \langle \phi_1, \dots, \phi_n \rangle$$

n times from \top .

Proof. Using the Bekic property.

Least fixed-points: splitting the roles of variables

Due to

$$\mu_x.\phi(x, x) = \mu_x.\mu_y.\phi(x, y)$$

Least fixed-points: splitting the roles of variables

Due to

$$\mu_x.\phi(x, x) = \mu_x.\mu_y.\phi(x, y)$$

we can separate computing the least fixed-points w.r.t:

weakly negative variables: variables that appear within the left-hand-side of an implication, from

fully positive variables: those appearing only within the right-hand-side of an implication.

Weakly negative least fixed-points: an example

Use

$$\mu_x.(f \circ g)(x) = f(\mu_y.(g \circ f)(y))$$

Weakly negative least fixed-points: an example

Use

$$\mu_x.(f \circ g)(x) = f(\nu_y.(g \circ f)(y))$$

to argue that:

$$\begin{aligned}\mu_x.[(x \rightarrow a) \rightarrow b] &= [\nu_y.(y \rightarrow b) \rightarrow a] \rightarrow b \\ &= [(\top \rightarrow b) \rightarrow a] \rightarrow b \\ &= [b \rightarrow a] \rightarrow b.\end{aligned}$$

Weakly negative least fixed-points: reducing to greatest fixed-points

If each occurrence of x in ϕ is weakly negative, then

$$\phi(x) = \phi_0[\phi_1(x)/y_1, \dots, \phi_n(x)/y_n]$$

with $\phi_0(y_1, \dots, y_n)$ negative in each y_j .

Due to

$$\mu_x.(f \circ g)(x) = f(\mu_y.(g \circ f)(y))$$

Weakly negative least fixed-points: reducing to greatest fixed-points

If each occurrence of x in ϕ is weakly negative, then

$$\phi(x) = \phi_0[\phi_1(x)/y_1, \dots, \phi_n(x)/y_n]$$

with $\phi_0(y_1, \dots, y_n)$ negative in each y_j .

Due to

$$\mu_x.(f \circ g)(x) = f(\mu_y.(g \circ f)(y))$$

we have

$$\begin{aligned}\mu_x.\phi(x) &= \mu_x.(\phi_0 \circ \langle \phi_1, \dots, \phi_n \rangle)(x) \\ &= \phi_0(\nu_{y_1 \dots y_n}.(\langle \phi_1, \dots, \phi_n \rangle \circ \phi_0)(y_1, \dots, y_n)).\end{aligned}$$

Interlude: least fixed-points of strong functions

If f and f_i , $i \in I$, are strong, then

$$\mu_x.a \wedge f(x) = a \wedge \mu_x.f(x),$$

$$\mu_x.\bigwedge_{i \in I} f_i(x) = \bigwedge_{i \in I} \mu_x.f_i(x),$$

$$\mu_x.a \rightarrow f(x) = a \rightarrow \mu_x.f(x).$$

Strongly positive fixed-points: disjunctive formulas

The equation

$$\mu_x. \bigwedge_{i \in I} f_i(x) = \bigwedge_{i \in I} \mu_x. f_i(x)$$

allows to push least fixed-points down through conjunctions.

Strongly positive fixed-points: disjunctive formulas

The equation

$$\mu_x. \bigwedge_{i \in I} f_i(x) = \bigwedge_{i \in I} \mu_x. f_i(x)$$

allows to push least fixed-points down through conjunctions.

Once all conjunctions have been pushed up in formulas, we are left to compute least fixed-points of *disjunctive formulas*, generated by the grammar:

$$\phi = x \mid \beta \vee \phi \mid \alpha \rightarrow \phi \mid \bigvee_{i=1, \dots, n} \phi_i,$$

where α and β do not contain the variable x .

We call α an *head subformula* and β a *side subformula*.

Least fixed-points of inflating functions

All functions f denoted by such formula ϕ are (monotone and) *inflating*:

$$x \leq f(x).$$

Let f_i , $i = 1, \dots, n$, be a collection of monotone inflating functions.
Then

$$\mu_x. \bigvee_{i=1, \dots, n} f_i(x) = \mu_x. (f_1 \circ \dots \circ f_n)(x).$$

Least fixed-points of disjunctive formulas

Proposition. Let ϕ be a disjunctive formula, with $Head(\phi)$ (resp., $Side(\phi)$) the collection of its head (resp., side) subformulas. Then

$$\mu_x.\phi = \bigwedge_{\alpha \in Head(\phi)} \alpha \rightarrow \bigvee_{\beta \in Side(\phi)} \beta.$$

Least fixed-points of disjunctive formulas

Proposition. Let ϕ be a disjunctive formula, with $Head(\phi)$ (resp., $Side(\phi)$) the collection of its head (resp., side) subformulas. Then

$$\mu_x.\phi = \bigwedge_{\alpha \in Head(\phi)} \alpha \rightarrow \bigvee_{\beta \in Side(\phi)} \beta.$$

If $Head(\phi) = \{\alpha_1, \dots, \alpha_n\}$ and $Side(\phi) = \{\beta_1, \dots, \beta_m\}$:

$$\begin{aligned} \mu_x.\phi &= \mu_x.\alpha_1 \rightarrow \alpha_2 \rightarrow \dots \rightarrow \alpha_n \rightarrow \beta_1 \vee \dots \vee \beta_m \vee x \\ &= \mu_x.\bigwedge_{\alpha \in Head(\phi)} \alpha \rightarrow \bigvee_{\beta \in Side(\phi)} \beta \vee x \\ &= \bigwedge_{\alpha \in Head(\phi)} \alpha \rightarrow \mu_x.\bigvee_{\beta \in Side(\phi)} \beta \vee x \\ &= \bigwedge_{\alpha \in Head(\phi)} \alpha \rightarrow \bigvee_{\beta \in Side(\phi)} \beta. \end{aligned}$$

Plan

A primer on mu-calculi

The intuitionistic μ -calculus

The elimination procedure

Bounding closure ordinals

Closure ordinals

Definition. (*Closure ordinal*). For \mathcal{K} a class of models and $\phi(x)$ a monotone formula/term, let

$$\text{cl}_{\mathcal{K}}(\phi) = \text{least ordinal } \alpha \text{ such that } \mathcal{M} \models \mu_x.\phi = \phi^\alpha(\perp).$$

In general, $\text{cl}_{\mathcal{K}}(\phi)$ might not exist.

If \mathcal{H} is the class of Heyting algebras and $\phi(x)$ is an intuitionistic formula, then

$$\text{cl}_{\mathcal{H}}(\phi) < \omega.$$

Upper bounds from fixed-point equations

$$\text{cl}_{\mathcal{K}}(f \circ g) \leq \text{cl}_{\mathcal{K}}(g \circ f) + 1,$$

$$\text{cl}_{\mathcal{H}}(\phi_0(\phi_1(x), \dots, \phi_n(x))) \leq n + 1,$$

when $\phi_0(y_1, \dots, y_n)$ contravariant,

$$\text{cl}_{\mathcal{H}}(\phi) \leq \mathbf{card}(\text{Head}(\phi)) + 1,$$

when ϕ is a disjunctive formula,

$$\text{cl}_{\mathcal{K}}(f \circ \Delta) \leq n \text{cl}_{\mathcal{K}}(g),$$

when $n = \text{cl}_{\mathcal{K}}(f(x, -))$ and $g(x) = \mu_y.f(x, y)$,

$$\text{cl}_{\mathcal{K}}(f \wedge g) \leq \text{cl}_{\mathcal{K}}(f) + \text{cl}_{\mathcal{H}}(g) - 1,$$

when f and g are strong.

Quizzes

1.

$$\bigwedge_{i=1, \dots, n} (x \rightarrow a_i) \rightarrow b_i$$

converges after k steps. $k = \dots?$

Quizzes

1.

$$\bigwedge_{i=1, \dots, n} (x \rightarrow a_i) \rightarrow b_i$$

converges after k steps. $k = \dots?$

2.

$$\bigvee_{i=1, \dots, n} (x \rightarrow a_i) \rightarrow b_i$$

converges after k steps. $k = \dots?$

After thoughts

- ▶ The elimination procedure gives a decision procedure for the Intuitionistic μ -calculus.
- ▶ A quadratic upper bound on the closure ordinals of intuitionistic formulas (yet, this does not yield polynomial equi-expressivity).
- ▶ General theory of fixed-point elimination: no uniform upper bounds, relevance of strongness.

References I



Alberucci, L. and Facchini, A. (2009).

The modal μ -calculus hierarchy on restricted classes of transition systems.
The Journal of Symbolic Logic, 74(4):1367–1400.



Bradfield, J. C. (1998).

The modal μ -calculus alternation hierarchy is strict.
Theor. Comput. Sci., 195(2):133–153.



Frittella, S. and Santocanale, L. (2014).

Fixed-point theory in the varieties \mathcal{D}_n .

In Höfner, P., Jipsen, P., Kahl, W., and Müller, M. E., editors, *Relational and Algebraic Methods in Computer Science - 14th International Conference, RAMiCS 2014, Marienstatt, Germany, April 28-May 1, 2014. Proceedings*, volume 8428 of *Lecture Notes in Computer Science*, pages 446–462. Springer.



Ghilardi, S., Gouveia, M. J., and Santocanale, L. (2016).

Fixed-point elimination in the intuitionistic propositional calculus.

In Jacobs, B. and Löding, C., editors, *Foundations of Software Science and Computation Structures - 19th International Conference, FOSSACS 2016, Held as Part of the European Joint Conferences on Theory and Practice of Software, ETAPS 2016, Eindhoven, The Netherlands, April 2-8, 2016, Proceedings*, volume 9634 of *Lecture Notes in Computer Science*, pages 126–141. Springer.



Ghilardi, S. and Zawadowski, M. W. (1997).

Model completions, r-Heyting categories.
Ann. Pure Appl. Logic, 88(1):27–46.



Kozen, D. (1983).

Results on the propositional mu-calculus.
Theor. Comput. Sci., 27:333–354.

References II



Lenzi, G. (1996).

A hierarchy theorem for the μ -calculus.

In auf der Heide, F. M. and Monien, B., editors, *Automata, Languages and Programming, 23rd International Colloquium, ICALP96, Paderborn, Germany, 8-12 July 1996, Proceedings*, volume 1099 of *Lecture Notes in Computer Science*, pages 87–97. Springer.



Pitts, A. M. (1992).

On an interpretation of second order quantification in first order intuitionistic propositional logic.

J. Symb. Log., 57(1):33–52.



Ruitenburg, W. (1984).

On the period of sequences ($a^n(p)$) in intuitionistic propositional calculus.

The Journal of Symbolic Logic, 49(3):892–899.



Santocanale, L. (2002).

The alternation hierarchy for the theory of μ -lattices .

Theory and Applications of Categories, 9:166–197.